

## Short Communication

# An ATP-Dependent Step Is Required for the Translocation of Microinjected Precursor mRNA into Nuclear Speckles

(ATP-dependent step / HeLa cell nucleus / microinjected splicing-competent RNA / RNA translocation into nuclear speckles)

V. KOPSKÝ<sup>1,2,3,+</sup>, J. VEČEŘOVÁ<sup>1,2,+</sup>, I. MELČÁK<sup>1,2,4</sup>, A. PLISS<sup>1,2</sup>, J. ŠTULÍK<sup>2</sup>,  
K. KOBERNA<sup>1,2</sup>, L. TOMÁŠKOVÁ<sup>1,2</sup>, I. RAŠKA<sup>1,2</sup>

<sup>1</sup>Department of Cell Biology, Institute of Experimental Medicine, Academy of Sciences of the Czech Republic, Prague, Czech Republic

<sup>2</sup>Laboratory of Gene Expression, 1<sup>st</sup> Medical Faculty, Charles University, Prague, Czech Republic

**Abstract.** Nuclear speckles (speckles) represent a distinct nuclear compartment within the interchromatin space and are enriched in splicing factors. In a previous study (Melčák et al., 2001), it has been shown that the pre-spliceosomal assembly on microinjected splicing-competent precursor mRNA takes place in the speckles, and it has been suggested that the targeting of RNA into speckles consists of two interdependent steps, namely the diffusion process, followed by the energy-dependent translocation of RNA into the speckles. In the present study, we confirm the existence of these two steps and show that this latter translocation is ATP dependent.

Most primary transcripts of mammalian protein-coding genes contain introns and have to be spliced before being transported to the cytoplasm. Precursor mRNA (pre-mRNA) splicing takes place in a large ribonucleoprotein complex termed spliceosome. The spliceosomes are generated by the constitutive assembly of U1, U2, U5, U4/U6 small nuclear ribonucleoprotein particles

(snRNPs) and various non-snRNP factors on pre-mRNAs in a cascade of sequence-specific steps (Steitz et al., 1988; Lamm and Lamond, 1993; Moore et al., 1993; Newman, 1994; Krämer, 1996). The formation of the functional spliceosome is thus preceded by the formation of a number of pre-spliceosomal complexes which contain, together with unspliced pre-mRNA, defined combinations of snRNP particles and non-snRNP factors (e.g. Steitz et al., 1988; Lamm and Lamond, 1993; Moore et al., 1993; Newman, 1994; Krämer, 1996).

When RNA polymerase II transcriptional and splicing components are mapped within the cell nucleus, they exhibit frequent sites of high local accumulation in the form of microclusters, which likely represent the sites of active transcription and co-transcriptional splicing (Neugebauer and Roth, 1997; Misteli and Spector, 1998). On the other hand, beside these microclusters, an accumulation of factors of the splicing apparatus is typically mapped to spatially distinct and large domains termed „speckles“, „SC35 domains“, the antibody to splicing factor SC35 (Fu and Maniatis, 1990) being used regularly for the visualization of speckles, or „splicing factor compartments“ (reviewed in Spector, 1990; Fakan, 1994; Misteli, 2000). Even though there is a consensus that speckles play a role in RNA metabolism, their exact function is presently unknown. These structures are not usually correlated with RNA polymerase II transcription, but it has been shown at the level of activation of some specific unique genes that speckles serve as pools of splicing factors, which are redistributed to the transcription/splicing sites (Misteli et al., 1997).

The splicing of pre-mRNAs may be a co-transcriptional event (Beyer and Osheim, 1988; Neugebauer and Roth, 1997; Custodio et al., 1999), and it has been shown that transcription and splicing are coupled through interactions of certain factors participating in both these processes (Corden and Patrajan, 1997; McCracken et al., 1997; Steinmetz,

Received January 11, 2002. Accepted January 28, 2002.

This work was supported by grants from the Grant Agency of the Czech Republic 304/00/1481 and 304/01/0729, from the Academy of Sciences of the Czech Republic IAA5039103 and AV0Z5039906, and from the Ministry of Education, Youth and Sports MSM: 111100003.


Corresponding author: Ivan Raška, Department of Cell Biology, Institute of Experimental Medicine, Academy of Sciences of the Czech Republic, Albertov 4, 128 00 Prague, Czech Republic, Tel: +420 (2) 24910315, Fax: +420 (2) 24917418, e-mail: iraska@lf1.cuni.cz.

<sup>3</sup>Present address: Department of Animal Physiology and Developmental Biology, Faculty of Science, Charles University, Viničná 7, Prague 2, 128 44, Czech Republic

<sup>4</sup>Present address: Laboratory of Cell Biology, Howard Hughes Medical Institute, The Rockefeller University, 1230 York Avenue, New York, NY 10021-6399, USA

+ V. K and J. V. were equally contributing authors

Abbreviations: pre-mRNA – precursor messenger RNA, snRNP – small nuclear ribonucleoprotein.



*Fig. 1.* (A, B) After the consecutive microinjection of apyrase (B) or buffer (A), and of the Ad1 RNA labelled with fluorochrome (in green), the cells were incubated at 37°C for 15 min. The cells were then labelled for the SC35 splicing factor (in red). The form of the Ad1 RNA, consisting of two exons and one intron, is shown in A. Note a part of a non-microinjected nucleus in A. The two left columns correspond to individual colour channels, the right column is the overlay. In A, RNA was accumulated within the speckles. In B, the ATP depletion resulted in the accumulation of RNA in dots adjacent to the speckles. In the quantitative evaluation (inserts in A and B), fluorescence intensities along the blue lines shown in A and B were scaled to the minimum and maximum values. The identical position of peaks A (insert) in the two colour channels testifies to the accumulation of RNAs in the speckles. Note the differences in the position of peaks in B (insert). The bar in A and B corresponds to 4  $\mu\text{m}$ .

1997). However, not all pre-mRNA sequences are processed co-transcriptionally and post-transcriptional splicing does occur (Zachar et al., 1993; Baurén and Wieslander, 1994; Wuarin and Schibler, 1994). Importantly, isolated pre-mRNAs from mammalian cells may contain both introns and poly(A) tails. Splicing may then be a post-transcriptional event, at least in some cases (McCracken et al., 1997; Minvielle-Sebastia and Keller, 1999), and it has been suggested that speckles are involved in post-transcriptional splicing (Melčák and Raška, 1996). With regard to the spatial relationship of pre-mRNA accumulations relative to speckle domains of splicing factor accumulations, primary transcripts of certain genes as well as the spliced RNAs have been mapped at sites of active transcription and/or outside the speckles (Zhang et al., 1994; Smith et al., 1999). However, pre-mRNAs from some other genes have been shown to be associated, or overlapped, with nuclear speckles (Xing et al., 1993, 1995; Huang and Spector, 1996; Dirks et al., 1997; Ishov et al., 1997; Jolly et al., 1999; Smith et al., 1999; Snaar et al., 1999; Johnson et al., 2000; Melčák et al., 2000).

The suggested function of speckles in splicing has been probed using a completely different approach in which the behavior of microinjected (and fluorochrome-labelled) pre-mRNA into the nuclei of HeLa cells was

investigated (Wang et al., 1991; Melčák et al., 2001). The rationale for this approach is based on the findings of Graessmann and Graessmann (1982) that microinjected intron-containing RNA is processed into functional mRNA within the cell nucleus. These exogenous RNAs skip the transcription context and behave similarly to endogenous unspliced RNAs released from the sites of transcription. Using model RNAs with a single intron, it has been shown that these RNAs rapidly accumulate in the speckles after microinjection in a process that is dependent on the intron (Wang et al., 1991; Melčák et al., 2001) and that the pre-spliceosome assembly on RNAs takes place within the speckles (Wang et al., 1991; Melčák et al., 2001). RNA targeting to and accumulation within the speckles is the result of the cumulative loading of splicing factors to the pre-mRNA.

In this study, we expand the results with microinjected adenovirus pre-mRNAs and demonstrate that the targeting of microinjected pre-mRNA, together with formed pre-spliceosomal complexes, to the speckles consists of two interdependent steps, namely the movement of pre-mRNA towards the speckles and its translocation into the speckles, this second step requiring ATP.

## Material and Methods

Synthesis of the fluorochrome-labelled and splicing-competent adenovirus Ad1 RNA, containing a single

intron, as well as HeLa cell culture, microinjection, SC35 immunolabelling and microscopy were performed according to Melčák et al. (2001).

In order to determine the effect of energy depletion on RNA localization, an ATP depletion by microinjected apyrase was performed. In order to reach this aim, consecutive microinjections were performed. Cells were microinjected with 500 U/ml of apyrase in 5 mM Tris-acetate, pH 6.95, and incubated for 5 min at 37°C. The same cells were then microinjected with RNA, washed with fresh prewarmed medium and incubated for 15 min at 37°C. The control cells were, instead of apyrase, microinjected with buffer only.

Four series of experiments were performed and 30 to 40 cells were microinjected in each experiment. However, not all microinjected cells could be subsequently investigated because a few cells either detached from the support or were disrupted. Also for this reason the described fluorescence pattern after apyrase microinjection (Fig. 1) was observed in 75, 85, 89 and 90% of microinjected cells.

## Results and Discussion

The control experiment was found to be compatible with the findings in previous studies (Wang et al., 1991; Melčák et al., 2001) and documented the rapid movement of microinjected splicing-competent Ad1 pre-mRNA into the speckles. The RNA was localized within the speckles, which were depicted by SC35 labelling (Fig. 1A).

A striking change in the RNA distribution was seen when the RNA microinjection was preceded by apyrase microinjection. RNA was also targeted towards the speckles, but remained accumulated outside this nuclear compartment. It formed dots of accumulated RNA adjacent to the speckles (Fig. 1B). Moreover, the speckles rounded up. As the ATP depletion necessarily led to transcription inhibition, the rounding up of the speckles was in agreement with previous results (e.g. O'Keefe et al., 1994; Melčák et al., 2000) documenting that the normal pattern of speckles is converted into round speckles in transcriptionally inactive cells.

The fluorescence pattern testified to the existence of an ATP-dependent step, which allowed for RNA accumulation within the speckles. Importantly, the fluorescence pattern was compatible with the experiment in which microinjected HeLa cells with Ad1 RNA were kept at 4°C, and strikingly resembled the fluorescence pattern observed with several microinjected mutant RNAs (Melčák et al., 2001). The mutant RNAs allow for the generation of pre-spliceosomal complexes, but only of such complexes which do not require ATP for their formation in *in vitro* experiments (Seraphin and Rosbach, 1989; Bennett et al., 1992; Michaud and Reed, 1993; Query et al., 1997; see Melčák et al. (2001) for detailed discussion).

In our previous study (Melčák et al., 2001), it was suggested that the targeting of microinjected splicing-competent RNA into speckles consisted of two interdependent steps, namely the movement of RNA towards the speckles, probably as a diffusion process, followed by the energy-dependent translocation of RNA into the speckles. The result of the performed ATP depletion experiment, which resulted, on the one hand, in the blocking of accumulation of the splicing-competent RNAs in the speckles, but, on the other, in their accumulation in the fluorescent dots adjacent to the speckles, demonstrates the existence of these two targeting steps, and supports the concept that nuclear speckles are involved in splicing.

## Acknowledgement

The authors wish to thank Dr. X.-D. Fu (Univ. California San Diego) for the anti-SC35 antibody, and Dr. C. Will (Univ. of Marburg) for the plasmid used for the preparation of RNA. They thank Dr. J. Malínský for help with the preparation of Figure and Dr. F. Raška for the English revision.

## References

- Baurén, G., Wieslander, L. (1994) Splicing of Balbiani ring 1 gene pre-mRNA occurs simultaneously with transcription. *Cell* **76**, 183-192.
- Bennett, M., Michaud, S., Kingston, J., Reed, R. (1992) Protein components specifically associated with pre-spliceosome and spliceosome complexes. *Genes Dev.* **6**, 1986-2000.
- Beyer, A. L., Osheim, Y. N. (1988) Splice site selection, rate of splicing, and alternative splicing on nascent transcripts. *Genes Dev.* **2**, 754-765.
- Corden, J. L., Patturajan, M. (1997) A CTD function linking transcription to splicing. *Trends Biochem. Sci.* **22**, 413-416.
- Custodio, N., Carmo-Fonseca, M., Geraghty, F., Pereira, H., Grosveld, F., Antoniou, M. (1999) Inefficient processing impairs release of RNA from the site of transcription. *EMBO J.* **18**, 2855-2866.
- Dirks, R. W., de Pauw, E. S. D., 97a Splicing factors associate with nuclear HCMV-IE transcripts after transcriptional activation of the gene, but dissociate upon transcription inhibition: evidence for a dynamic organization of splicing factors. *J. Cell Sci.* **110**, 505-513.
- Fakan, S. (1994) Perichromatin fibrils are in situ forms of nascent transcripts. *Trends Cell. Biol.* **4**, 86-90.
- Fu, X. D., Maniatis, T. (1990) Factor required for mammalian spliceosome assembly is localized to discrete regions in the nucleus. *Nature* **343**, 437-441.
- Graessmann, M., Graessmann, A. (1982) Simian virus 40 cRNA is processed into functional mRNA in microinjected monkey cells. *EMBO J.* **1**, 1081-1088.
- Huang, S., Spector, D. L. (1996) Intron-dependent recruitment of pre-mRNA splicing factors to sites of transcription. *J. Cell Biol.* **133**, 719-732.
- Ishov, A. M., Stenberg, R. M., Maul, G. G. (1997) Human cytomegalovirus immediate early interaction with host nuclear structures: definition of an immediate transcript environment. *J. Cell Biol.* **138**, 5-16.

- Johnson, C., Primorac, D., McKinstry, M., McNeil, J., Rowe, D., Lawrence, J. B. (2000) Tracking COL1A1 RNA in osteogenesis imperfecta: splice-defective transcripts initiate transport from the gene but are retained within the SC35 domain. *J. Cell Biol.* **150**, 417-431.
- Jolly, C., Vourc'h, C., Robert-Nicoud, M., Morimoto, R. I. (1999). Intron-dependent association of splicing factors with active genes. *J. Cell Biol.* **145**, 1133-1143
- Krämer, A. (1996) The structure and function of proteins involved in mammalian pre-mRNA splicing. *Annu. Rev. Biochem.* **65**, 367-409.
- Lamm, G. M., Lamond, A. I. (1993) Non-snRNP protein splicing factors. *Biochim. Biophys. Acta* **1173**, 247-265.
- McCracken, S., Fong, N., Yankulov, K., Ballantyne, S., Pan, G., Greenblatt, J., Patterson, S. D., Wickens, M., Bentley, D. L. (1997) The C-terminal domain of RNA polymerase II couples mRNA processing to transcription. *Nature* **385**, 357-361.
- Melčák, I., Raška, I. (1996) Structural organization of the pre-mRNA splicing commitment. A hypothesis. *J. Struct. Biol.* **117**, 189-194.
- Melčák, I., Cermanová, Š., Jirsová, K., Koberna, K., Malinský, J., Raška, I. (2000) Nuclear pre-mRNA compartmentalization: trafficking of released transcripts to splicing factor reservoirs. *Mol. Biol. Cell* **11**, 497-510.
- Melčák, I., Melčáková, Š., Kopský, V., Večeřová, J., Raška, I. (2001) Prespliceosomal assembly on microinjected precursor mRNA takes place in nuclear speckles. *Mol. Biol. Cell* **12**, 393-406.
- Michaud, S., Reed, R. (1993) A functional association between the 5' and 3' splice site is established in the earliest prespliceosome complex (E) in mammals. *Genes Dev.* **7**, 1008-1020.
- Minvielle-Sebastia, L., Keller, W. (1999) mRNA polyadenylation and its coupling to other RNA processing reactions and to transcription. *Curr. Opin. Cell Biol.* **11**, 352-357.
- Misteli, T. (2000) Cell biology of transcription and pre-mRNA splicing: nuclear architecture meets nuclear function. *J. Cell Sci.* **113**, 1841-1849.
- Misteli, T., Spector, D. L. (1998) RNA polymerase II targets pre-mRNA splicing factors to transcription sites in vivo. *Mol. Cell* **3**, 697-705.
- Misteli, T., Cáceres, J. F., Spector, D. L. (1997) The dynamics of a pre-mRNA splicing factor in living cells. *Nature* **387**, 523-527.
- Moore, M. J., Query, C. C., Sharp, P. A. (1993) Splicing of precursors to messenger RNAs by the spliceosome. In: *The RNA World*, eds. Gesteland, R. F., Atkins, J. F., pp. 303-357, CSHL Press, Cold Spring Harbor.
- Neugebauer, K. M., Roth, M. B. (1997) Distribution of pre-mRNA splicing factors at sites of RNA polymerase II transcription. *Genes Dev.* **11**, 1148-1159.
- Newman, A. J. (1994) Small nuclear RNAs and pre-mRNA splicing. *Curr. Opin. Cell Biol.* **6**, 360-367.
- O'Keefe, R. T., Mayeda, A., Sadowski, C. L., Krainer, A. R., Spector, R. L. (1994) Disruption of pre-mRNA splicing in vivo results in reorganization of splicing factors. *J. Cell Biol.* **124**, 249-260.
- Query, C. C., McCaw, P. S., Sharp, P. A. (1997) A minimal spliceosomal complex A recognizes the branch site and polypyrimidine tract. *Mol. Cell Biol.* **17**, 2944-2953.
- Seraphin, B., Rosbash, M. (1989) Identification of functional U1 snRNA-pre-mRNA complexes committed to spliceosome assembly and splicing. *Cell* **59**, 349-358.
- Smith, K. P., Moen, P. T., Wydner, K. L., Coleman, J. R., Lawrence, J. B. (1999) Processing of endogenous pre-mRNAs in association with SC-35 domains is gene specific. *J. Cell Biol.* **144**, 617-629.
- Snaar, S. P., Vincent, M., Dirks, R. W. (1999) RNA polymerase II localizes at sites of human cytomegalovirus immediate-early RNA synthesis and processing. *J. Histochem. Cytochem.* **47**, 245-254.
- Spector, D. L. (1990) Higher order nuclear organization: three-dimensional distribution of small nuclear ribonucleoprotein particles. *Proc. Natl. Acad. Sci. USA* **87**, 147-151.
- Steinmetz, E. J. (1997) Pre-mRNA processing and the CTD of RNA polymerase II: the tail that wags the dog? *Cell* **89**, 491-494.
- Steitz, J. A., Black, D. A., Gerke, V., Parker, K. A., Krämer, A., Frendewey, D., Keller, W. (1988) Functions of the abundant U-snRNPs. In: *Structure and Function of Major and Minor Small Nuclear Ribonucleoprotein Particles*, ed. Birnstiel, M. L., pp. 71-79, Springer-Verlag, Berlin.
- Wang, J., Cao, L. G., Wang, Y. L., Pederson, T. (1991) Localization of pre-messenger RNA at discrete nuclear sites. *Proc. Natl. Acad. Sci. USA* **88**, 7391-7395.
- Wuarin, J., Schibler, U. (1994) Physical isolation of nascent RNA chains transcribed by RNA polymerase II: evidence for cotranscriptional splicing. *Mol. Cell Biol.* **14**, 7219-7225.
- Xing, Y., Johnson, C. V., Dobner, P., Lawrence J. B. (1993) Higher level organization of individual gene transcription and RNA splicing: integration of nuclear structure and function. *Science* **259**, 1326-1330.
- Xing, Y., Johnson, C. V., Moen, P. T. Jr., McNeil, J. A., Lawrence, J. B. (1995) Nonrandom gene organization: structural arrangements of specific pre-mRNA transcription and splicing with SC-35 domains. *J. Cell Biol.* **131**, 1635-1647.
- Zachar, Z., Kramer, J., Mims, I. P., Bingham, P. M. (1993) Evidence for channeled diffusion of pre-mRNAs during nuclear RNA transport in metazoans. *J. Cell Biol.* **121**, 729-742.
- Zhang, G., Taneja, K. L., Singer, R. H., Green, M. R. (1994) Localization of pre-mRNA splicing in mammalian nuclei. *Nature* **372**, 809-812.